## **Prescribing Information**

## **AnhydroI™ Forte**

20% w/v Cutaneous Solution

Aluminium chloride hexahydrate 20% w/v.

**Uses:** For the topical treatment of hyperhidrosis specifically involving axillae, hands or feet.



**Directions:** Adults, children and the elderly: Apply to the affected sites at night, as required, and allow to dry. Wash off in the morning.

Contra-indications, warnings, side effects etc: Please refer to SPC for full details before prescribing. Do not use if sensitive to any of the ingredients. Care should be taken to restrict the application to the affected sites only. Do not bathe or shower immediately before use and, if the axillae are treated, do not shave or use depilatories on this area within 12 hours before or after use. If applied too frequently, Anhydrol Forte may cause irritation which should be treated with a mild topical hydrocortisone cream. Keep away from the eyes and mucous membranes. Care should be taken to avoid Anhydrol Forte coming into direct contact with clothing, polished surfaces, jewellery or metal. Highly flammable: keep away from flames.

Package quantity, NHS price and MA number: 60ml roll-on bottle. £2.99, PL 00173/0030

Legal category: P.

**MA holder**: Dermal Laboratories Ltd, Tatmore Place, Gosmore, Hitchin,

Herts SG4 7QR, UK.

**Date of preparation:** September 2025.

'Anhydrol' is a trademark.

Adverse events should be reported. Reporting forms and information can be found at yellowcard.mhra.gov.uk Adverse events should also be reported to Dermal.

To find out more visit: https://www.dermal.co.uk/healthcare-professionals/products/anhydrol-forte/

